

CONSTRUCTING ALGEBRAIC CLOSURES

KEITH CONRAD

Let K be a field. We want to construct an algebraic closure of K , *i.e.*, an algebraic extension of K which is algebraically closed. It will be built as the quotient of a polynomial ring in a very large number of variables.

For each nonconstant monic polynomial $f(X)$ in $K[X]$, let its degree be n_f and let $t_{f,1}, \dots, t_{f,n_f}$ be independent variables. Let $A = K[\{t_{f,i}\}]$ be the polynomial ring generated over K by independent variables doubly indexed by every nonconstant monic $f \in K[X]$ and $1 \leq i \leq n_f$. This is a *very large* polynomial ring containing K .

Let I be the ideal in A generated by the coefficients of all the differences

$$f(X) - \prod_{i=1}^{n_f} (X - t_{f,i}) \in A[X]$$

as f runs over nonconstant monic polynomials in $K[X]$. Working modulo I we have $f(X) \equiv \prod_i (X - t_{f,i})$, so $f(X)$ splits into linear factors in $(A/I)[X]$. We want to use a maximal ideal in place of I since working modulo a maximal ideal would give a complete splitting of every $f(X)$ from $K[X]$ over a *field*.

Lemma 1. *The ideal I is proper: $1 \notin I$.*

Proof. This will follow from the existence of a splitting field for finitely many polynomials in $K[X]$.

Suppose $1 \in I$. Then we can write 1 as a finite sum $\sum_{j=1}^m a_j c_j$, where each c_j is a coefficient in some difference

$$(1) \quad f_j(X) - \prod_{i=1}^{n_j} (X - t_{f_j,i})$$

and $a_j \in A$. Here $f_j(X)$ is monic in $K[X]$ and $n_j = \deg(f_j)$. There is a (finite) field extension L/K in which the finitely many f_j 's all split completely, say $f_j(X) = \prod_{i=1}^{n_j} (X - r_{j,i})$ in $L[X]$. (Some numbers in the list $r_{j,1}, \dots, r_{j,n_j}$ might be repeated.) We can use the roots $r_{j,i}$ of $f_1(X), \dots, f_m(X)$ to construct a ring homomorphism φ from $A = K[\{t_{f,i}\}]$ to L by substitution: φ fixes K , $\varphi(t_{f_j,i}) = r_{j,i}$ for $1 \leq i \leq n_j$, and $\varphi(t_{f,i}) = 0$ if f is not one of the f_j 's. Extend φ to a homomorphism $A[X] \rightarrow L[X]$ by acting on coefficients. The polynomial in (1) is mapped by φ to

$$f_j(X) - \prod_{i=1}^{n_j} (X - r_{f_j,i}) = 0 \text{ in } L[X],$$

so every coefficient in (1) is mapped by φ to 0 in L . In particular, $\varphi(c_j) = 0$. Thus φ sends the equation $1 = \sum_{j=1}^m a_j c_j$ in A to the equation $1 = 0$ in L , and that is a contradiction. \square

Since I is a proper ideal, Zorn's lemma guarantees that I is contained in some maximal ideal \mathfrak{m} in A . (I suspect I itself is not a maximal ideal, but I don't have a proof of that.)

The quotient ring $A/\mathfrak{m} = K[\{t_{f,i}\}]/\mathfrak{m}$ is a field and the natural composite homomorphism $K \rightarrow A \rightarrow A/\mathfrak{m}$ of rings let us view the field A/\mathfrak{m} as an extension of K (ring homomorphisms out of fields are always injective).

Theorem 2. *The field A/\mathfrak{m} is an algebraic closure of K .*

Proof. For a nonconstant monic $f(X) \in K[X]$ we have $f(X) - \prod_{i=1}^{n_f} (X - t_{f,i}) \in I[X] \subset \mathfrak{m}[X]$, so in $(A/\mathfrak{m})[X]$ we have $f(X) = \prod_i (X - \bar{t}_{f,i})$, where $\bar{t}_{f,i}$ denotes $t_{f,i} \bmod \mathfrak{m}$. Each $\bar{t}_{f,i}$ is algebraic over K (being a root of $f(X)$) and A is generated as a ring over K by the $t_{f,i}$'s, so A/\mathfrak{m} is generated as a ring over K by the $\bar{t}_{f,i}$'s. Therefore A/\mathfrak{m} is an algebraic extension field of K in which every nonconstant monic in $K[X]$ splits completely.

We will now show A/\mathfrak{m} is algebraically closed, and thus it is an algebraic closure of K . Set $F = A/\mathfrak{m}$. It suffices to show every monic irreducible $\pi(X)$ in $F[X]$ has a root in F . We have already seen that each nonconstant monic polynomial in $K[X]$ splits completely in $F[X]$, so let's show $\pi(X)$ is a factor of some monic polynomial in $K[X]$. There is a root α of $\pi(X)$ in some extension of F . Since α is algebraic over F and F is algebraic over K , α is algebraic over K . That implies some monic $f(X)$ in $K[X]$ has α as a root. The polynomial $\pi(X)$ is the minimal polynomial of α in $F[X]$, so $\pi(X) \mid f(X)$ in $F[X]$. Since $f(X)$ splits completely in $F[X]$, $\alpha \in F$. \square

Our construction of an algebraic closure of K is done, but we want to compare it with another construction to put the one above in context.

The idea of building an algebraic closure of K by starting with a large polynomial ring over K whose variables are indexed by polynomials in $K[X]$ goes back at least to Emil Artin. He used a large polynomial ring (somewhat smaller than the ring $K[\{t_{f,i}\}]$ we started with above) modulo a suitable maximal ideal to obtain an algebraic extension K_1/K such that every nonconstant polynomial in $K[X]$ has a root in K_1 (not, *a priori*, that they all split completely in $K_1[X]$). Then he iterated this construction with K_1 in place of K to get a new algebraic extension K_2/K_1 , and so on, and proved that the union $\bigcup_{n \geq 1} K_n$ (or, more rigorously, the direct limit of the K_n 's) *contains* an algebraic closure of K [2, pp. 544-545]. With more work, treating separately characteristic 0 and characteristic p , it can be shown [3] that Artin's construction only needs one step: K_1 is an algebraic closure of K (so $K_n = K_1$ for all n , which is not obvious in Artin's own proof). In other words, the following is true: if F/K is an algebraic extension such that every nonconstant polynomial in $K[X]$ has a root in F then every nonconstant polynomial in $F[X]$ has a root in F , so F is an algebraic closure of K .

Theorem 2 and its proof, which I learned from B. Conrad, is a variation on a proof by Zorn [6] in the paper where he first introduced Zorn's lemma. It modifies Artin's construction by using a larger polynomial ring over K in order to adjoin to K in one step a full set of roots – not just one root – of each nonconstant monic in $K[X]$, rather than one root for each polynomial. This makes it easier to prove the constructed field is an algebraic closure of K . A similar construction, using a maximal ideal in a tensor product, is in [1, Prop. 4, p. A V 21].

Remark 3. That every field has an algebraic closure and that two algebraic closures of a field are isomorphic were first proved by Steinitz in 1910 in a long paper that created from scratch the general theory of fields as part of abstract algebra. The influence of this paper on the development of algebra was enormous; for an indication of this, see [4]. Steinitz was hindered in this work by the primitive state of set theory at that time and he used the

well-ordering principle rather than Zorn's lemma (which only became widely known in the 1930s [6]). Steinitz's proof of the existence of algebraic closures and their uniqueness up to isomorphism, together with his account of set theory, took up 20 pages [5, Sect. 19–21].

At the end of the proof of Theorem 2, the polynomial $f(X)$ in $K[X]$ with α as a root can be taken to be irreducible over K , so we could build an algebraic closure of K by defining the ideal I using just the monic *irreducible* $f(X)$ in $K[X]$ rather than all monic $f(X)$ in $K[X]$; the proofs of Lemma 1 and Theorem 2 carry over with no essential changes other than inserting the word “irreducible” in a few places. Finally, if we restrict the f in the construction of I to run over the monic separable polynomials in $K[X]$, or the monic separable irreducible polynomials in $K[X]$, then the field A/\mathfrak{m} turns out to be a separable closure of K . The proof of Lemma 1 carries over with the f_j being separable (or separable irreducible), and in the proof of Theorem 2 two changes are needed: A/\mathfrak{m} is a separable algebraic extension of K since it would be generated as a ring over K by roots of separable polynomials in $K[X]$, and we need transitivity of separability instead of algebraicity (if F/K is separable algebraic then each root of a separable polynomial in $F[X]$ is separable over K).

REFERENCES

- [1] N. Bourbaki, “Algebra II: Chapters 4-7,” Springer-Verlag, New York, 1990.
- [2] D. Dummit, R. Foote, “Abstract Algebra,” 3rd ed., Wiley, New York, 2004.
- [3] R. Gilmer, *A Note on the Algebraic Closure of a Field*, Amer. Math. Monthly **75** (1968), 1101-1102.
- [4] P. Roquette, *In memoriam Ernst Steinitz (1871–1928)*, J. Reine Angew. Math. **648** (2010), 1–11. Online at <https://www.mathi.uni-heidelberg.de/~roquette/STEINITZ.pdf>.
- [5] E. Steinitz, *Algebraische Theorie der Körper*, J. Reine Angew. Math. **137** (1910), 167–309. Online at <https://eudml.org/doc/149323>.
- [6] M. Zorn, *A Remark on a Method in Transfinite Algebra*, Bull. Amer. Math. Society **41** (1935), 667–670.