

THE p -ADIC GROWTH OF HARMONIC SUMS

KEITH CONRAD

The numbers

$$H_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n} = \sum_{k=1}^n \frac{1}{k}$$

are called *harmonic sums*. Integrals can approximate these sums as real numbers, *e.g.*, H_n and $\int_1^n dt/t = \log n$ differ by a bounded amount ($H_n - \int_1^n dt/t$ is positive and monotonically decreasing), so the H_n 's slowly diverge in \mathbf{R} . We are interested here in the arithmetic behavior of harmonic sums, *i.e.*, their p -adic growth. This is definitely not an important topic, but it is interesting to compare what happens with this rational sequence across the real and different p -adic landscapes.

Here are some initial values:

n	H_n
1	1
2	3/2
3	11/6
4	25/12
5	137/60
6	49/20
7	363/140
8	761/280
9	7129/2520
10	7381/2520
11	83711/27720

TABLE 1. Harmonic sums

For $n > 1$, it appears that $H_n \notin \mathbf{Z}$. More precisely, the numerator of H_n is odd and the denominator of H_n is even. This suggests a strategy for proving H_n is not an integer.

Theorem 1 (Theisinger). *For all $n \geq 2$, $H_n \notin \mathbf{Z}$.*

Proof. Let L be the least common multiple of $1, 2, \dots, n$, so H_n can be written as a fraction with denominator L . For $1 \leq k \leq n$, write $L = ka_k$ with $a_k \in \mathbf{Z}^+$, so $1/k = a_k/L$. Then

$$H_n = \sum_{k=1}^n \frac{1}{k} = \frac{\sum_{k=1}^n a_k}{L}.$$

Since $n \geq 2$, L is even. We will show $\sum_{k=1}^n a_k$ is odd, so the ratio is not integral.

Set 2^r to be the largest power of 2 up to n : $2^r \leq n < 2^{r+1}$. The only integer up to n divisible by 2^r is 2^r itself, since $2 \cdot 2^r > n$. Therefore $L = 2^r b$ where b is odd, so $2^r b = ka_k$ for $1 \leq k \leq n$. When $k = 2^r$ we see that $a_k = b$ is odd. When $k \neq 2^r$, k is not divisible by 2^r , so a_k must be even. Therefore in the numerator $\sum_{k=1}^n a_k$, one term (at $k = 2^r$) is odd and the rest are even, so the total sum is odd. \square

Theisinger [4] did not prove Theorem 1 using anything p -adic: he used determinants and Bertrand's postulate, which says there is always a prime between n and $2n$ when $n \geq 2$.¹

The proof above for Theorem 1 presents it as a 2-adic result: $H_n \notin \mathbf{Z}$ because $H_n \notin \mathbf{Z}_2$. Here is a second proof of Theorem 1 brings out 2-adic features more clearly.

Proof. Let $2^r \leq n < 2^{r+1}$, so $r \geq 1$ and the highest power of 2 that appears in some reciprocal in the sum defining H_n is 2^r . The only reciprocal in H_n with denominator divisible by 2^r is $1/2^r$. Indeed, every other such reciprocal would be $1/(2^r c)$ for odd $c > 1$, but if that term is in the sum then so is $1/(2^r \cdot 2) = 1/2^{r+1}$, which is false since $2^{r+1} > n$. Therefore $1/2^r$ has a more highly negative 2-adic valuation than every other term in H_n , so it is not cancelled out in the sum. This means $H_n \notin \mathbf{Z}_2$, so $H_n \notin \mathbf{Z}$. \square

This proof gives a formula for the 2-adic valuation of the harmonic sums: $\text{ord}_2(H_n) = -r$ where $2^r \leq n < 2^{r+1}$. To see this formula in action, we rewrite the initial harmonic sums in Table 2 with the power of 2 in the denominators made explicit. The exponent jumps when n is a power of 2.

n	H_n
1	1
2	$3/2$
3	$11/2 \cdot 3$
4	$25/2^2 \cdot 3$
5	$137/2^2 \cdot 15$
6	$49/2^2 \cdot 5$
7	$363/2^2 \cdot 35$
8	$761/2^3 \cdot 35$
9	$7129/2^3 \cdot 315$
10	$7381/2^3 \cdot 2520$
11	$83711/2^3 \cdot 3465$

TABLE 2. Harmonic sums viewed 2-adically

Since $|H_n|_2 = 2^r$ if $2^r \leq n < 2^{r+1}$, $|H_n|_2 \rightarrow \infty$ as $n \rightarrow \infty$ (explicitly, $n/2 < |H_n|_2 \leq n$). Does $|H_n|_p \rightarrow \infty$ as $n \rightarrow \infty$ for odd primes p ? We will address this later.

Theorem 2 (Kürschák). *For $m \leq n - 2$, $H_n - H_m \notin \mathbf{Z}_2$. In particular, $H_n - H_m \notin \mathbf{Z}$.*

Taking $m = 1$ and $n \geq 3$ recovers Theorem 1 for $n \geq 3$ since $H_1 \in \mathbf{Z}$. Theorem 2 is false if $m = n - 1$ and n is odd, since then $H_n - H_m = 1/n \in \mathbf{Z}_2$.

Proof. Writing

$$H_n - H_m = \sum_{k=m+1}^n \frac{1}{k},$$

we will show there is a unique term in the sum with the most negative 2-adic valuation. Let $r = \max_{m < k \leq n} \text{ord}_2(k)$. Since $n \geq m + 2$, the sum $H_n - H_m$ has at least two terms in it, so some k is even and therefore $r \geq 1$.

We will show there is only one integer from m up to $n - 1$ with 2-adic valuation r . Suppose there are two such numbers. Write them as $2^r c$ and $2^r d$ with odd $c < d$. Then $c + 1$ is even and $2^r c < 2^r(c + 1) < 2^r d$, so $1/(2^r(c + 1))$ appears in $H_n - H_m$. But $\text{ord}_2(2^r(c + 1)) \geq r + 1$ since c is odd. This contradicts the definition of r . Therefore there is only one term in $H_n - H_m$ with 2-adic valuation $-r$, so $\text{ord}_2(H_n - H_m) = -r$. \square

¹Theisinger referred to Bertrand's postulate as Chebyshev's theorem, since Chebyshev had proved it.

This type of 2-adic argument (in the setting of rational numbers, not 2-adic numbers) is due to Kürschák [3], so he should be credited with the 2-adic proof of Theorem 1.

We now turn to the p -adic behavior of H_n for $p \neq 2$. The material below is based on the notes “Harmonics and Primes” by Nicholas Rogers.

The harmonic sums H_1, H_2, \dots, H_{p-1} are all p -adically integral, since the denominators appearing in them are prime to p and thus are p -adic integers (even p -adic units). Since $1/p$ is outside the ring \mathbf{Z}_p , $H_p = H_{p-1} + 1/p$ is outside \mathbf{Z}_p : $\text{ord}_p(H_p) = -1$. It need not be true (as it is for $p = 2$) that $H_n \notin \mathbf{Z}_p$ for all $n \geq p$. For example, H_3, H_4 and H_5 are outside \mathbf{Z}_3 (all with 3-adic valuation -1), but H_6, H_7 , and H_8 are in \mathbf{Z}_3 . Then $H_9, H_{10}, \dots, H_{20}$ are outside of \mathbf{Z}_3 . The harmonic sums return to \mathbf{Z}_3 for H_{21}, H_{22} , and H_{23} , then leave \mathbf{Z}_3 and come back in again for H_{66}, H_{67} , and H_{68} . Then the harmonic sums leave \mathbf{Z}_3 and never return. In fact, $|H_n|_3 \rightarrow \infty$ as $n \rightarrow \infty$. We will not discuss the proof of this, but the next theorem gives a simple connection between p -adic integrality and p -adic divergence of the harmonic sums.

Theorem 3. *For a prime p , the following conditions are equivalent:*

- (1) $\{n \geq 1 : H_n \in \mathbf{Z}_p\}$ is finite,
- (2) $|H_n|_p \rightarrow \infty$ as $n \rightarrow \infty$.

Proof. Trivially the second condition implies the first. To prove the (more interesting) reverse implication, pick $n \geq 1$ and write $n = pq + r$ with $0 \leq r \leq p - 1$. Since $1/k \in \mathbf{Z}_p$ when $(p, k) = 1$, in the quotient group $\mathbf{Q}_p/\mathbf{Z}_p$ a harmonic sum is equal to the sum of its terms whose denominators are divisible by p , so

$$H_n \equiv \frac{1}{p} + \frac{1}{2p} + \dots + \frac{1}{pq} = \frac{1}{p}H_q,$$

so $H_n - (1/p)H_q \in \mathbf{Z}_p$.

We are assuming that $|H_n|_p \leq 1$ only finitely many times, so there is some $N_0 \geq 1$ such that $|H_n|_p > 1$ for $n \geq N_0$. We will show this N_0 controls the p -adic divergence of the harmonic sums: for all $k \geq 0$,

$$n \geq p^k N_0 \implies |H_n|_p > p^k.$$

For $k = 0$ this is true by the definition of N_0 . Assuming it is true for k , suppose $n \geq p^{k+1}N_0$. Writing $n = pq + r$ with $0 \leq r \leq p - 1$, from $pq + r \geq p^{k+1}N_0$ we must have $q \geq p^k N_0$, so $|H_q|_p > p^k$. Therefore $|(1/p)H_q|_p > p^{k+1}$. Since $H_n - (1/p)H_q \in \mathbf{Z}_p$,

$$\left| H_n - \frac{1}{p}H_q \right|_p \leq 1 < p^{k+1} = \left| \frac{1}{p}H_q \right|_p,$$

so by the nonarchimedean property $|H_n|_p = |(1/p)H_q|_p > p^{k+1}$. □

The same reasoning shows $|H_n|_p \rightarrow \infty$ as $n \rightarrow \infty$ if and only if the set $\{n \geq 1 : H_n \in p\mathbf{Z}_p\}$ is finite. (Replace the strict inequalities $|H_n|_p > 1$ and $|H_n|_p > p^k$ in the proof with $|H_n|_p \geq 1$ and $|H_n|_p \geq p^k$.) This set, introduced in [2], is denoted

$$J(p) = \{n \geq 1 : H_n \in p\mathbf{Z}_p\}.$$

To say $n \notin J(p)$ means H_n has no p in its numerator. For instance, Theorem 1 says $J(2) = \emptyset$. In [2] it is proved that $J(3) = \{2, 7, 22\}$, $J(5) = \{4, 20, 24\}$, and

$$J(7) = \{6, 42, 48, 295, 299, 337, 341, 2096, 2390, 14675, 16731, 16735, 102728\}.$$

The set $J(11)$ has 638 elements [1]. It is conjectured that $J(p)$ is finite for all p , which by Theorem 3 is equivalent to saying $|H_n|_p \rightarrow \infty$ as $n \rightarrow \infty$ for all p .

The rest of our discussion will focus on the particular sum H_{p-1} .

Theorem 4. For each odd prime p , $H_{p-1} \in p\mathbf{Z}_p$. Therefore $J(p) \neq \emptyset$ for $p > 2$.

Proof. In the sum

$$H_{p-1} = 1 + \frac{1}{2} + \cdots + \frac{1}{p-1},$$

each term is a unit in \mathbf{Z}_p , so we can reduce modulo p and check that $H_{p-1} \equiv 0 \pmod{p\mathbf{Z}_p}$. The integers $\{1, 2, \dots, p-1\}$ represent the units modulo p , so their inverses do as well. Thus, after replacing each $1/k$ with the integer from 1 to $p-1$ that is equal to it in $\mathbf{Z}_p/(p)$,

$$\begin{aligned} H_{p-1} &\equiv 1 + 2 + \cdots + p-1 \pmod{p} \\ &\equiv \frac{p(p-1)}{2} \pmod{p} \\ &\equiv 0 \pmod{p}, \end{aligned}$$

where in the last step we needed $p \neq 2$. □

In Table 3, we factor the initial harmonic sums into primes, and can see in this data (for $p = 5, 7$, and 11) that not only is H_{p-1} divisible by p , but by p^2 .

n	H_n
1	1
2	$3/2$
3	$11/2 \cdot 3$
4	$5^2/2^2 \cdot 3$
5	$137/2^2 \cdot 3 \cdot 5$
6	$7^2/2^2 \cdot 5$
7	$3 \cdot 11^2/2^2 \cdot 5 \cdot 7$
8	$761/2^3 \cdot 5 \cdot 7$
9	$7129/2^3 \cdot 3^2 \cdot 5 \cdot 7$
10	$11^2 \cdot 61/2^3 \cdot 3^2 \cdot 5 \cdot 7$
11	$97 \cdot 863/2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$

TABLE 3. Harmonic sums fully factored

Theorem 5 (Wolstenholme [5], 1862). For each prime $p \geq 5$, $H_{p-1} \in p^2\mathbf{Z}_p$.

Proof. We collect terms in H_{p-1} that are equidistant from the middle of the sum:

$$\begin{aligned} H_{p-1} &= 1 + \frac{1}{2} + \cdots + \frac{1}{p-1} \\ &= \left(1 + \frac{1}{p-1}\right) + \left(\frac{1}{2} + \frac{1}{p-2}\right) + \cdots + \left(\frac{1}{(p-1)/2} + \frac{1}{(p+1)/2}\right) \\ &= \sum_{k=1}^{(p-1)/2} \left(\frac{1}{k} + \frac{1}{p-k}\right) \\ &= \sum_{k=1}^{(p-1)/2} \frac{p}{k(p-k)} \\ &= p \sum_{k=1}^{(p-1)/2} \frac{1}{k(p-k)}. \end{aligned}$$

Since a p has been pulled out, we want to show this last sum is in $p\mathbf{Z}_p$. The terms in the sum are p -adic units, and reducing the terms modulo p yields

$$\sum_{k=1}^{(p-1)/2} \frac{1}{k(p-k)} \equiv - \sum_{k=1}^{(p-1)/2} \frac{1}{k^2} \pmod{p\mathbf{Z}_p}.$$

The numbers $1^2, \dots, ((p-1)/2)^2$ represent all the nonzero squares modulo p , so their reciprocals also represent the nonzero squares modulo p . Therefore

$$\sum_{k=1}^{(p-1)/2} \frac{1}{k^2} \equiv \sum_{k=1}^{(p-1)/2} k^2 \pmod{p\mathbf{Z}_p}.$$

Using the formula $\sum_{k=1}^n k^2 = n(n+1)(2n+1)/6$ with $n = (p-1)/2$,

$$\sum_{k=1}^{(p-1)/2} k^2 = \frac{p(p^2-1)}{24}.$$

Since $p > 3$, $(p, 24) = 1$, so this sum is in $p\mathbf{Z}_p$ and we're done. \square

Theorem 5 says $\text{ord}_p(H_{p-1}) \geq 2$. If one calculates H_{p-1} for primes $3 < p < 10000$, always $\text{ord}_p(H_{p-1}) = 2$. But at $p = 16843$ and at $p = 2124679$, $\text{ord}_p(H_{p-1}) = 3$. No further examples where $\text{ord}_p(H_{p-1}) > 2$ are known.

REFERENCES

- [1] D. W. Boyd, *A p -adic study of the partial sums of the harmonic series*, Experiment. Math **3** (1994), 287–302.
- [2] A. Eswarathasan and E. Levine, *p -integral harmonic sums*, Discrete Math. **91** (1991), 249–257.
- [3] J. Kürschák, *A Harmonikus Sorról*, Mat. és Fiz. Lapok, **27** (1918), 299–300. Online at http://real-j.mtak.hu/7278/1/MTA_MatematikaiEsPhysikaiLapok_27.pdf.
- [4] L. Theisinger *Bemerkung über die harmonische Reihe*, Montash. f. Math. und Physik **26** (1915), 132–134. Online at <http://www.literature.at/viewer.alo?objid=12428&viewmode=fullscreen&page=136>.
- [5] J. Wolstenholme, *On certain properties of prime numbers*, The Quarterly Journal of Pure and Applied Mathematics **5** (1862), 35–39. Online at <https://babel.hathitrust.org/cgi/pt?id=hvd.32044102924370&view=1up&seq=53>.