

COSETS AND LAGRANGE'S THEOREM

KEITH CONRAD

1. INTRODUCTION

Pick an integer $m \neq 0$. For $a \in \mathbf{Z}$, the congruence class $a \bmod m$ is the set of integers $a + mk$ as k runs over \mathbf{Z} . We can write this set as $a + m\mathbf{Z}$. This can be thought of as a translated subgroup: start with the subgroup $m\mathbf{Z}$ and add a to it. This idea can be carried over from \mathbf{Z} to an arbitrary group, provided we distinguish between translation of a subgroup on the left and on the right.

Definition 1.1. Let G be a group and H be a subgroup. For $g \in G$, the sets

$$gH = \{gh : h \in H\}, \quad Hg = \{hg : h \in H\}$$

are called, respectively, a *left H -coset* and a *right H -coset*.

In other words, a coset is what we get when we take a subgroup and shift it: a left coset gH is shifted on the left side and a right coset Hg is shifted on the right side.¹ The best way to think about cosets is that they are *shifted subgroups* or *translated subgroups*.

Note g lies in both gH and Hg , since $g = ge = eg$. Typically $gH \neq Hg$. When G is abelian, though, left and right cosets of a subgroup by a common element are the same. When an abelian group operation is written additively, an H -coset should be written as $g + H$, which is the same as $H + g$.

Example 1.2. In the additive group \mathbf{Z} , with subgroup $m\mathbf{Z}$, the $m\mathbf{Z}$ -coset of a is $a + m\mathbf{Z}$. This is just a congruence class modulo m .

Example 1.3. In the group \mathbf{R}^\times , with subgroup $H = \{\pm 1\}$, the H -coset of x is $xH = \{x, -x\}$. This is “ x up to sign.”

Example 1.4. When $G = S_3$, and $H = \{(1), (12)\}$, the table below lists the left H -cosets and right H -cosets of every element of the group. Compute a few of them for non-identity elements to satisfy yourself that you understand how they are found.

g	gH	Hg
(1)	{(1), (12)}	{(1), (12)}
(12)	{(1), (12)}	{(1), (12)}
(13)	{(13), (123)}	{(13), (132)}
(23)	{(23), (132)}	{(23), (123)}
(123)	{(13), (123)}	{(23), (123)}
(132)	{(23), (132)}	{(13), (132)}

Notice first of all that cosets are usually not subgroups (some do not even contain the identity). Also, since $(13)H \neq H(13)$, a particular element can have different left and right H -cosets. Since $(13)H = (123)H$, different elements can have the same left H -coset. (You have already seen this happen with congruences: $14 + 3\mathbf{Z} = 2 + 3\mathbf{Z}$, since $14 \equiv 2 \pmod{3}$.)

¹Marshall Hall's book *The Theory of Groups* calls gH a right coset and Hg a left coset. Don't do that.

In the next section we will see how cosets look in some geometric examples, where we can visualize cosets. Then we will see that cosets arise in certain decimal patterns and as “inhomogeneous solution spaces” to linear equations or differential equations. Then we will look at some general properties of cosets. The *index* of a subgroup in a group, which tells us how many cosets the subgroup has (either on the right or on the left), will lead to the most basic important theorem about finite groups: Lagrange’s theorem. We will see a few applications of Lagrange’s theorem and finish up with the more abstract topics of left and right coset spaces and double coset spaces.

2. GEOMETRIC EXAMPLES OF COSETS

When a group is defined in terms of vectors and matrices, we can often get a picture of the group and its cosets.

Example 2.1. Let $G = \mathbf{R}^2$ and $H = \mathbf{R}e_1$ be the x -axis. The (left) H -coset of a vector $\mathbf{v} \in \mathbf{R}^2$ is

$$\mathbf{v} + H = \mathbf{v} + \mathbf{R}e_1 = \{\mathbf{v} + ce_1 : c \in \mathbf{R}\}.$$

This is the line parallel to H (the x -axis) that passes through the endpoint of \mathbf{v} . The H -cosets in general are the lines parallel to H . Two parallel lines are either equal or disjoint, so each pair of H -cosets is either equal or disjoint. In Figure 1, the H -cosets of \mathbf{v} and \mathbf{v}' are equal while those of \mathbf{v} and \mathbf{w} are disjoint.




FIGURE 1. The cosets of $\mathbf{R}e_1$ in \mathbf{R}^2 .

Example 2.2. Let $G = \text{Aff}^+(\mathbf{R})$, the 2×2 matrices $\begin{pmatrix} x & y \\ 0 & 1 \end{pmatrix}$ with $x > 0$ under matrix multiplication. Geometrically, we identify elements of G with points (x, y) in the plane where $x > 0$. Such points form a right half-plane:

$$\begin{pmatrix} x & y \\ 0 & 1 \end{pmatrix} \longleftrightarrow (x, y) \in \mathbf{R}_{>0} \times \mathbf{R}.$$

The identity element becomes the point $(1,0)$. See Figure 2. This is quite similar to the idea of identifying complex numbers $x + yi$ with points (x, y) in the plane.




FIGURE 2. The group $\text{Aff}^+(\mathbf{R})$.

Unlike the geometric interpretation of complex number addition, which is ordinary vector addition with the intuitive “parallelogram rule”, the group law on $\text{Aff}^+(\mathbf{R})$ when written in terms of points in the plane is unfamiliar: starting from

$$\begin{pmatrix} x & y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} u & v \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} xu & xv + y \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} x & y \\ 0 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1/x & -y/x \\ 0 & 1 \end{pmatrix},$$

we get the group law on points

$$(x, y)(u, v) = (xu, xv + y), \quad (x, y)^{-1} = \left(\frac{1}{x}, -\frac{y}{x}\right).$$

Each matrix $\begin{pmatrix} x & y \\ 0 & 1 \end{pmatrix}$ in $\text{Aff}^+(\mathbf{R})$ breaks up as the product $\begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$ (not as $\begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix}$!). These special types of matrices each form subgroups, which we will call H and K :

$$(2.1) \quad H = \left\{ \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} : y \in \mathbf{R} \right\}, \quad K = \left\{ \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} : x > 0 \right\}.$$

They are pictured in Figure 3.


For $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ in $\text{Aff}^+(\mathbf{R})$, what are its left and right H -cosets $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}H$ and $H\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$? To compute these multiply a general element of H on the left and right by $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$:

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} a & ay + b \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} a & b + y \\ 0 & 1 \end{pmatrix}.$$

Here a and b are fixed, while y varies over \mathbf{R} . The numbers $ay + b$ run over all of \mathbf{R} and the numbers $b + y$ run over \mathbf{R} . This means the left and right H -cosets of $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ in $\text{Aff}^+(\mathbf{R})$ are the same:

$$(2.2) \quad \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} H = H \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} = \left\{ \begin{pmatrix} a & t \\ 0 & 1 \end{pmatrix} : t \in \mathbf{R} \right\}.$$

In the picture of $\text{Aff}^+(\mathbf{R})$, this is simply the vertical line parallel to H passing through the point (a, b) . See the first picture in Figure 4.

FIGURE 3. The group $\text{Aff}^+(\mathbf{R})$.

To compute the left K -coset of $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$, for $x > 0$ we have

$$(2.3) \quad \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} ax & b \\ 0 & 1 \end{pmatrix}.$$

As $x > 0$ varies, the right side of (2.3) runs through all matrices of the form $\begin{pmatrix} t & b \\ 0 & 1 \end{pmatrix}$ with $t > 0$. Therefore

$$(2.4) \quad \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} K = \left\{ \begin{pmatrix} t & b \\ 0 & 1 \end{pmatrix} : t > 0 \right\}.$$

In particular, the left K -coset of $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ is determined by b alone and is independent of the choice of $a > 0$: $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} K = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} K$. This is the horizontal line through (a, b) . See the second picture in Figure 4.

To find out what the right K -coset of $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ is, multiply a general element of K by $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ on the right:

$$\begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} ax & bx \\ 0 & 1 \end{pmatrix}.$$

Letting $x > 0$ vary, we obtain

$$(2.5) \quad K \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} = \left\{ \begin{pmatrix} u & v \\ 0 & 1 \end{pmatrix} : u > 0, v = \frac{b}{a}u \right\}.$$

This is a half-line out of the origin with slope b/a (third picture in Figure 4). In particular, the left and right K -cosets of $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ in $\text{Aff}^+(\mathbf{R})$ are *not* the same if $b \neq 0$. A right K -coset of $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ depends on both a and b , or more precisely on their ratio b/a , while its left K -coset depends only on b .

For both H and K , its collection of all left cosets and its collection of all right cosets fill up the group $\text{Aff}^+(\mathbf{R})$ in Figure 5 *without overlap*: a family of parallel half-lines, either horizontal or vertical, and a family of half-lines out of the origin.

Later in this handout, we will see that left (or right) cosets of a subgroup in a group exhibit properties that can be seen in our geometric examples: different left cosets of a




FIGURE 4. Left and Right Cosets of H and K in $\text{Aff}^+(\mathbf{R})$.




FIGURE 5. Left and Right Coset Decompositions of $\text{Aff}^+(\mathbf{R})$ by H and K .

subgroup are disjoint, and the collection of all left cosets of a subgroup cover the group (likewise for right cosets).

3. COSETS AND DECIMAL EXPANSIONS

Each rational number a/b whose denominator b is relatively prime to 10 has a decimal expansion that is purely periodic. For example,

$$\frac{1}{3} = .333333\dots, \quad \frac{1}{7} = .142857142857142857\dots, \quad \frac{15}{41} = .609756097560975\dots$$

where the repeating part has 1 term, 6 terms, and 5 terms respectively. We abbreviate the decimal expansion by writing only the periodic part and put a line over it, so

$$\frac{1}{3} = .\overline{3}, \quad \frac{1}{7} = .\overline{142857}, \quad \frac{15}{41} = .\overline{60975}.$$

In the tables below are all the *reduced* proper fractions with denominator 7, 13, and 27. (We don't include $3/27$ in the second table, for instance, since it is not reduced.)

Fraction	Decimal	Fraction	Decimal	Fraction	Decimal
1/7	$\overline{.142857}$	1/13	$\overline{.076923}$	2/13	$\overline{.153846}$
2/7	$\overline{.285714}$	3/13	$\overline{.230769}$	4/13	$\overline{.307692}$
3/7	$\overline{.428571}$	5/13	$\overline{.384615}$	6/13	$\overline{.461538}$
4/7	$\overline{.571428}$	7/13	$\overline{.538461}$	8/13	$\overline{.615384}$
5/7	$\overline{.714285}$	9/13	$\overline{.692307}$	10/13	$\overline{.769230}$
6/7	$\overline{.857142}$	11/13	$\overline{.846153}$	12/13	$\overline{.923076}$

Fraction	Decimal	Fraction	Decimal	Fraction	Decimal
1/27	$\overline{.037}$	2/27	$\overline{.074}$	4/27	$\overline{.148}$
5/27	$\overline{.185}$	7/27	$\overline{.259}$	8/27	$\overline{.296}$
10/27	$\overline{.370}$	11/27	$\overline{.407}$	13/27	$\overline{.481}$
14/27	$\overline{.518}$	16/27	$\overline{.592}$	17/27	$\overline{.629}$
19/27	$\overline{.703}$	20/27	$\overline{.740}$	22/27	$\overline{.814}$
23/27	$\overline{.851}$	25/27	$\overline{.925}$	26/27	$\overline{.962}$

There are two interesting features revealed in this table:

- the decimal expansions for all the *reduced* fractions with the same denominator have the same period length (denominator 7 has period length 6, denominator 13 has period length 6, and denominator 27 has period length 3),
- if we cyclically shift the digits in the repeating part of the decimal expansions of a reduced fraction, we get the repeating part of the decimal expansion of another reduced fraction with the same denominator.

The first feature is explained by Euler's theorem from elementary number theory, and we won't discuss it here.² We want to focus on the second feature. First, let's make sure we understand what it is saying. The string of digits 142857 is the repeating part of the decimal for 1/7, and if we shift the numbers one position to the left, getting 428571, we look in the first table and see it is the repeating part of the decimal for 3/7. If we shift one position to the left again, we get 285714, and when we look at the first table we find that is the repeating part of the decimal for 2/7. Similarly, 407 is the repeating part of the decimal for 11/27, and if we shift the terms one position to the left we get 074, which the second table tells us is the repeating part of the decimal for 2/27. Are you surprised by this? It turns out this phenomenon, which even a grade school student could discover, is actually cosets in disguise. This is best explained by an example.

Example 3.1. Consider 5/13, which the first table tells us is $\overline{.384615}$. Multiplying by 10 has the effect of cyclically shifting the digits in the decimal one position to the left:

$$10 \cdot \frac{5}{13} = 10(\overline{.384615}) = 10(.384615384615\dots) = 3.84615384615\dots = 3.\overline{846153}.$$

The numerator of $10(5/13)$ is 50, and $50 \equiv 11 \pmod{13}$, so $50/13 = 11/13 + \text{integer}$. Since we already saw that $50/13 = 3.\overline{846153}$, the integer part of $50/13$ is 3 and therefore $11/13 = \overline{.846153}$. In other words, shifting the digits in $\overline{.384615}$ one position to the left turns 5/13 into 11/13, and the reason is that shifting digits in a decimal one position to the left corresponds to multiplication by 10 and $5 \cdot 10 \equiv 11 \pmod{13}$.

If we want to shift the digits to the left one more time, the fraction we get will again be obtained from multiplying the new numerator by 10 and reducing the product mod 13: $11 \cdot 10 = 110 \equiv 6 \pmod{13}$, or in terms of the original numerator, $5 \cdot 10^2 \equiv 6 \pmod{13}$. In

²See Theorem 4.3 in <https://kconrad.math.uconn.edu/blurbs/ugradnumthy/eulerthm.pdf>.

general, if $5 \cdot 10^k \equiv r \pmod{13}$, where $0 \leq r < 13$, then $r/13$ will have a decimal expansion whose repeating part is the digits for $5/13$ shifted to the left k times. All these decimals have the same length, which is 6, and that means after we do this shifting 6 times we come back to the original fraction and have run through all the cycle shifts of the digits in the decimal for $5/13$. The fractions we have created in this way are all $r/13$ where $r \equiv 5 \cdot 10^k \pmod{13}$ for some k . In other words, the numerators of the fractions with denominator 13 having a decimal with the same digit cycle (up to the choice of starting point) as the decimal for $5/13$ is precisely $\{5 \cdot 10^k \pmod{13} : k = 1, 2, \dots\}$, which is a coset in $(\mathbf{Z}/(13))^\times$ of the cyclic subgroup $\langle 10 \pmod{13} \rangle$.

In general, if a/b is a reduced fraction where $(10, b) = 1$, then the other reduced fractions whose decimal expansion has a repeating block that is a shift of the repeating block of the decimal for a/b are precisely those fractions with denominator b and numerator in the coset $a\langle 10 \pmod{b} \rangle$ of $(\mathbf{Z}/(b))^\times$.

Example 3.2. The fractions $2/27, 11/27$, and $20/27$ all have decimals with digit cycle 074 up to a shift, in $(\mathbf{Z}/(27))^\times$ we have $\langle 10 \pmod{27} \rangle = \{1, 10, 19\}$, and modulo 27 the numerators of the fractions are $\{2, 11, 20\} = \{2, 2 \cdot 10^2, 2 \cdot 10\} = 2\langle 10 \pmod{27} \rangle$, which is a coset of $\langle 10 \pmod{27} \rangle$ in $(\mathbf{Z}/(27))^\times$.

4. INHOMOGENEOUS SOLUTION SPACES ARE COSETS

When solving a linear equation $A\mathbf{x} = \mathbf{b}$ (A is a matrix), where the right side is non-zero, we call the equation inhomogeneous. The general solution to this type of equation is $\mathbf{x}_0 + \mathbf{y}$, where \mathbf{x}_0 is a *particular* solution of $A\mathbf{x} = \mathbf{b}$ and \mathbf{y} is the *general* solution to the (homogeneous) equation $A\mathbf{x} = \mathbf{0}$. Let's see how the solutions to an inhomogeneous linear equation or inhomogeneous linear differential equation are an example of a coset. I do not think the two examples that follow will give *insight* into cosets; they simply serve to show that cosets appear in courses usually taken before abstract algebra.

Example 4.1. Consider a system of linear equations, written as a matrix equation $A\mathbf{x} = \mathbf{b}$, where A is an $m \times n$ real matrix, $\mathbf{b} \in \mathbf{R}^m$, and \mathbf{x} is an unknown vector in \mathbf{R}^n . Let W be the solution set of the corresponding homogeneous equation, where the right side is $\mathbf{0}$:

$$W = \{\mathbf{x} \in \mathbf{R}^n : A\mathbf{x} = \mathbf{0}\}.$$

This is a group under vector addition. (It contains $\mathbf{0} \in \mathbf{R}^n$, if $A\mathbf{x} = \mathbf{0}$ and $A\mathbf{x}' = \mathbf{0}$ then $A(\mathbf{x} + \mathbf{x}') = A\mathbf{x} + A\mathbf{x}' = \mathbf{0} + \mathbf{0} = \mathbf{0}$, and $A(-\mathbf{x}) = -A(\mathbf{x}) = -\mathbf{0} = \mathbf{0}$.) For $\mathbf{b} \neq \mathbf{0}$, if the equation $A\mathbf{x} = \mathbf{b}$ has a solution, say \mathbf{x}_0 , then the general solution to $A\mathbf{x} = \mathbf{b}$ is $\mathbf{x}_0 + W$. Indeed, for each $\mathbf{x} \in \mathbf{R}^n$ we have the following equivalences:

$$\begin{aligned} A\mathbf{x} = \mathbf{b} &\iff A\mathbf{x} = A\mathbf{x}_0 \quad \text{since } A\mathbf{x}_0 = \mathbf{b} \\ &\iff A(\mathbf{x} - \mathbf{x}_0) = \mathbf{0} \\ &\iff \mathbf{x} - \mathbf{x}_0 \in W \\ &\iff \mathbf{x} \in \mathbf{x}_0 + W. \end{aligned}$$

The solution set $\mathbf{x}_0 + W$ is a coset of W in \mathbf{R}^n .

Remark 4.2. Consider the linear differential equation

$$(4.1) \quad y' + xy = x^3.$$

The solutions to $y' + xy = 0$, where the right side is 0 (homogeneous case), are the functions of the form $ce^{-x^2/2}$. A particular solution to the equation (4.1) is $x^2 - 2$, and anyone who had a good course in differential equations will know the general solution to (4.1) must take the form $x^2 - 2 + ce^{-x^2/2}$. Letting c vary, the solution set to (4.1) is $x^2 - 2 + \mathbf{R}e^{-x^2/2}$. This is a coset of functions, so the same phenomenon we met with linear equations also occurs for linear differential equations.

5. PROPERTIES OF COSETS

We will generally focus our attention on left cosets of a subgroup. Proofs of the corresponding properties of right cosets will be completely analogous, and can be worked out by the reader.

Since $g = ge$ lies in gH , every element of G lies in some left H -coset, namely the left coset defined by the element itself. (Take a look at Example 1.4, where (13) lies in $(13)H$.) Similarly, $g \in Hg$ since $g = eg$.

A subgroup is always a left and a right coset of itself: $H = eH = He$. (This is saying nothing other than the obvious fact that if we multiply all elements of a subgroup by the identity, on either the left or the right, we get nothing new.) What is more important to recognize is that we can have $gH = H$ (or $Hg = H$) even when g is not the identity. For instance, in the additive group \mathbf{Z} , $10 + 5\mathbf{Z} = 5\mathbf{Z}$. All this is saying is that if we shift the multiples of 5 by 10, we get back the multiples of 5. Isn't that obvious? In fact, the only way we can have $a + 5\mathbf{Z} = 5\mathbf{Z}$ is if a is a multiple of 5, *i.e.*, if $a \in 5\mathbf{Z}$.

For a subgroup H of a group G , and $g \in G$, when does gH equal H ?

Theorem 5.1. *For $g \in G$, $gH = H$ if and only if $g \in H$.*

Proof. Since $g = ge \in gH$, having $gH = H$ certainly requires $g \in H$.

Now we need to show that if $g \in H$, then $gH = H$. We prove $gH = H$ by showing each is a subset of the other. Since $g \in H$, $gh \in H$ for all $h \in H$, so $gH \subset H$. To see $H \subset gH$, note $h = g(g^{-1}h)$ and that $g^{-1}h$ is in H (since $g^{-1} \in H$). \square

Example 5.2. Consider the subgroup $H = \{1, s\}$ of D_4 . We have

$$sH = \{s, s^2\} = \{s, e\} = H.$$

Two of the elements of D_4 that are not in H are r and r^2 . Their left H -cosets are not equal to H :

$$rH = \{r, rs\}, \quad r^2H = \{r^2, r^2s\}.$$

Notice the left H -cosets H , rH , and r^2H are not just unequal (which means each has an element not in another), but are in fact *disjoint* (which means no element of one is in another). This is similar to Example 2.1, where the cosets are a family of parallel lines: different parallel lines are disjoint. This is a completely general phenomenon, as follows.

Theorem 5.3. *If two left H -cosets share a common element, then they are equal. Equivalently, two left H -cosets that are not equal have no elements in common, *i.e.*, they are disjoint.*

Proof. To show left H -cosets with a common element are the same, suppose x belongs to g_1H and to g_2H , say

$$(5.1) \quad x = g_1h_1 = g_2h_2$$

where $h_1, h_2 \in H$. Then $g_1 = g_2 h_2 h_1^{-1}$. Each element of $g_1 H$ has the form $g_1 h$ for some $h \in H$, and

$$g_1 h = g_2 (h_2 h_1^{-1} h) \in g_2 H.$$

Since h was arbitrary in H , we see $g_1 H \subset g_2 H$. The reverse inclusion, $g_2 H \subset g_1 H$, follows by a similar argument (use the equation $g_2 = g_1 h_1 h_2^{-1}$ instead). \square

By Theorem 5.3, no element lies in more than one left H -coset. We call an element of a left coset a *representative* of that coset. A set of representatives for all the left H -cosets is called a *complete set* of left coset representatives.

Example 5.4. By the table in Example 1.4, each element of S_3 is in exactly one of H , $(13)H$, and $(23)H$, so

$$(5.2) \quad S_3 = H \cup (13)H \cup (23)H.$$

This is a union of disjoint sets. An example of a complete set of left coset representatives of H is $(1), (13),$ and (23) . Another complete set of left coset representatives of H is $(12), (13),$ and (132) .

Example 5.5. Let $G = \mathbf{Z}$ and $H = m\mathbf{Z}$, for $m > 0$. The coset decomposition of H in G (left and right is the same) is just the decomposition of \mathbf{Z} into congruence classes modulo m :

$$(5.3) \quad \mathbf{Z} = m\mathbf{Z} \cup (1 + m\mathbf{Z}) \cup (2 + m\mathbf{Z}) \cup \cdots \cup (m - 1 + m\mathbf{Z}).$$

It is standard to use $\{0, 1, \dots, m - 1\}$ as the coset representatives.

Example 5.6. Let $G = \text{Aff}^+(\mathbf{R})$ and $K = \left\{ \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} : x > 0 \right\}$. We saw in Example 2.2 that a left K -coset of a matrix depends only on the upper-right entry of the matrix: $\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} K = \left\{ \begin{pmatrix} x & b \\ 0 & 1 \end{pmatrix} : x > 0 \right\}$. Moreover, since all matrices in a given left K -coset have a common upper-right entry, we can use the upper-right entry to parametrize different left K -cosets:

$$(5.4) \quad \text{Aff}^+(\mathbf{R}) = \bigcup_{b \in \mathbf{R}} \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} K,$$

where the union is disjoint. Coset representatives are the matrices $\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$.

Example 5.7. Consider the (additive) group \mathbf{R} and the subgroup \mathbf{Z} . Every real number, up to addition by an integer, looks like a number between 0 and 1 (more precisely, in the half-open interval $[0, 1)$). Different numbers in this range are not \mathbf{Z} -translates of each other, so $[0, 1)$ is a complete set of (left) coset representatives of \mathbf{Z} in \mathbf{R} :

$$(5.5) \quad \mathbf{R} = \bigcup_{0 \leq x < 1} (x + \mathbf{Z}),$$

where the sets making up this union are disjoint.

Example 5.8. We saw in Example 2.1, where $G = \mathbf{R}^2$, that the cosets of the x -axis $\mathbf{R}\mathbf{e}_1$ in the plane are the lines parallel to the x -axis. A complete set of left coset representatives of $\mathbf{R}\mathbf{e}_1$ is a choice of one point on each line parallel to the x -axis. These choices could be made at random, but a nicer method is to use the points lying on a line (any line) not parallel to the x -axis. Such a line will pass once through each line parallel to the x -axis. An example of such a line is the y -axis, *i.e.*, the y -axis is a set of coset representatives for $\mathbf{R}\mathbf{e}_1$ in \mathbf{R}^2 :

$$(5.6) \quad \mathbf{R}^2 = \bigcup_{y \in \mathbf{R}} ((0, y) + \mathbf{R}\mathbf{e}_1),$$

where the union is disjoint. Coset representatives are $\{(0, y) : y \in \mathbf{R}\}$.

The decompositions in (5.2), (5.3), (5.4), (5.5), and (5.6) have analogues in every group G . For a subgroup H of G , every element of G lies in some left H -coset, and different left H -cosets are disjoint, so we can write G as a union of disjoint left H -cosets. Write these different cosets as $g_i H$, so we have a disjoint union

$$(5.7) \quad G = \bigcup_{i \in I} g_i H.$$

Here I is just an indexing set counting the number of different left H -cosets in G . Formula (5.7) is called the left H -coset decomposition of G . The particular multipliers g_i (which are nothing other than a particular choice of one element from each left H -coset) are a complete set of left coset representatives of H in G .

In a group, the passage from H to gH by left multiplication by g leads to no collapsing of terms. For instance, in Example 5.4, the cosets of $\{(1), (12)\}$ in S_3 all have size 2. Even in cases of an infinite subgroup, as in Example 2.1, the cosets all look like the original subgroup (just shifted). Let's prove such a comparison between a subgroup and each of its cosets in general.

Theorem 5.9. *Let H be a subgroup of the group G . Each left H -coset in G has a bijection with H . In particular, when H is finite, the cosets of H all have the same size as H .*

Proof. Pick a left coset, say gH . We can pass from gH to H by left multiplication by g^{-1} : $g^{-1}(gh) = h \in H$. Conversely, we can pass from H to gH by left multiplication by g . These functions from gH to H and *vice versa* are inverses to each other, showing gH and H are in bijection with each other. \square

Of course there is an analogous result for right cosets, which the reader can formulate.

6. THE INDEX AND LAGRANGE'S THEOREM

For an integer $m \neq 0$, the number of cosets of $m\mathbf{Z}$ in \mathbf{Z} is $|m|$. This gives us an interesting way to think about the meaning of $|m|$, other than its definition as “ m made positive.” Passing from \mathbf{Z} to other groups, counting the number of cosets of a subgroup gives a useful numerical invariant.

Definition 6.1. Let H be a subgroup of a group G . The *index* of H in G is the number of left cosets of H in G . This number, which is a positive integer or ∞ , is denoted $[G : H]$.

The terminology “index” is due to Cauchy in 1815 to count something that turns out to be $[G : H]$ in a special case. Concretely, the index of a subgroup tells us how many times we have to translate the subgroup around (on the left) to cover the whole group.

Example 6.2. Since $H = \{(1), (12)\}$ has 3 left cosets in S_3 , by Example 5.4, $[S_3 : H] = 3$.

Example 6.3. The subgroup $H = \{1, s\}$ of D_4 has four left cosets:

$$H, \quad rH = \{r, rs\}, \quad r^2H = \{r^2, r^2s\}, \quad r^3H = \{r^3, r^3s\}.$$

The index of H in D_4 is 4.

Example 6.4. For a positive integer m , $[\mathbf{Z} : m\mathbf{Z}] = m$, since $0, 1, \dots, m-1$ are a complete set of coset representatives of $m\mathbf{Z}$ in \mathbf{Z} .

Example 6.5. What is the index of $15\mathbf{Z}$ inside $3\mathbf{Z}$? (Not inside \mathbf{Z} , but $3\mathbf{Z}$.) Modulo 15, a multiple of 3 is congruent to 0, 3, 6, 9, or 12. That is, we have the disjoint union

$$3\mathbf{Z} = 15\mathbf{Z} \cup (3 + 15\mathbf{Z}) \cup (6 + 15\mathbf{Z}) \cup (9 + 15\mathbf{Z}) \cup (12 + 15\mathbf{Z}).$$

Thus $[3\mathbf{Z} : 15\mathbf{Z}] = 5$.

Example 6.6. The index $[\mathbf{R} : \mathbf{Z}]$ is infinite, by Example 5.7: there are infinitely many cosets of \mathbf{Z} in \mathbf{R} .

Remark 6.7. If we allow ourselves to use the language of cardinal numbers, which permits a distinction between different orders of infinity, then we could define $[G : H]$ as the cardinality of the set of left H -cosets. This would have no effect on the meaning of a finite index, but would allow a more refined meaning in the case of an infinite index. Since we will not have much use for the index concept when it is infinite, we stick with the more concrete approach in Definition 6.1.

In the case of finite groups, there is a simple formula for the index of a subgroup.

Theorem 6.8. *When G is a finite group, and H is a subgroup, $[G : H] = |G|/|H|$.*

For example, this formula says the index of $\{(1), (12)\}$ in S_3 is $6/2 = 3$. Compare with the computation of the same index in Example 6.2. Theorem 6.8 lets us read off the index of the subgroup just from knowing the size of the subgroup, without actual coset constructions.

Proof. Since G is finite, H has finitely many left cosets in G . Let $t = [G : H]$, and write the different left cosets of H as g_1H, \dots, g_tH . We know that two left cosets of H are either the same or are disjoint. Therefore we have

$$(6.1) \quad G = g_1H \cup \dots \cup g_tH,$$

where the union is disjoint. By Theorem 5.9, each left H -coset has the same size as H , so computing the size of both sides of (6.1) tells us

$$(6.2) \quad |G| = t|H|.$$

Thus $[G : H] = t = |G|/|H|$. □

Using the formula for the index as a ratio, we get the next “transitivity” result quite easily in the case of finite groups.

Theorem 6.9. *In a group G , indices are multiplicative in towers: for subgroups $K \subset H \subset G$,*

$$[G : K] = [G : H][H : K]$$

if $[G : K]$ is finite or if $[G : H]$ and $[H : K]$ are finite. In particular, if G is finite then $[G : K] = [G : H][H : K]$.

An example of this when G is an infinite group is $G = \mathbf{Z}$, $H = 3\mathbf{Z}$, and $K = 15\mathbf{Z}$: $[G : K] = 15$, $[G : H] = 3$ and $[H : K] = 5$.

Proof. First suppose G is a finite group. Theorem 6.8 implies

$$[G : H][H : K] = \frac{|G|}{|H|} \cdot \frac{|H|}{|K|} = \frac{|G|}{|K|} = [G : K].$$

Now suppose G is possibly not finite. We will treat separately the cases when the right side of the desired formula is finite and the left side is finite.

Case 1: $[G : H]$ and $[H : K]$ are finite. Let $m = [G : H]$ and $n = [H : K]$. Then G is the union of m different left H -cosets g_1H, \dots, g_mH and H is the union of n different left K -cosets h_1K, \dots, h_nK . We want to show $[G : K] = mn$, *i.e.*, G is a union of mn different left K -cosets.

Since $G = \bigcup_{i=1}^m g_iH$ and $H = \bigcup_{j=1}^n h_jK$, we have $g_iH = \bigcup_{j=1}^n g_ih_jK$, so

$$G = \bigcup_{i=1}^m g_iH = \bigcup_{i=1}^m \bigcup_{j=1}^n g_ih_jK,$$

so $[G : K] \leq mn$. To prove this inequality is an equality, we show the cosets g_ih_jK are different as i and j vary.

Suppose $g_ih_jK = g_{i'}h_{j'}K$ for some i, j, i', j' . We will show $i = i'$ and $j = j'$. Since $K \subset H$, we get $g_ih_jK \subset g_iH$ and $g_{i'}h_{j'}K \subset g_{i'}H$, so g_iH and $g_{i'}H$ overlap. That means $g_iH = g_{i'}H$ (Theorem 5.3), so $i = i'$ (the terms g_1, \dots, g_m lie in different left H -cosets). Therefore $g_ih_jK = g_ih_{j'}K$, so $h_jK = h_{j'}K$. That implies $j = j'$ since h_1, \dots, h_n are in different K -cosets.

Case 2: $[G : K] < \infty$. Since $K \subset H \subset G$, left K -cosets in H are special cases of left K -cosets in G , so $[H : K] \leq [G : K] < \infty$. Each left H -coset in G is a union of left K -cosets in G , so from $[G : K]$ being finite we get $[G : H] \leq [G : K]$. Thus $[G : H]$ and $[H : K]$ are finite, so by Case 1, $[G : K] = [G : H][H : K]$. \square

In Theorem 6.8, where G is finite, we found that $|H|$ divides $|G|$ since the ratio $|G|/|H|$ is a positive integer. This is called Lagrange's theorem.

Theorem 6.10 (Lagrange, 1771). *In a finite group, the order of each subgroup divides the order of the group.*

Proof. Let G be a finite group, and H a subgroup. By (6.2), $|H|$ divides $|G|$. In fact, we found the ratio $|G|/|H|$ counts the number of left H -cosets in G . \square

We will see some applications of Lagrange's theorem in the next section.

The converse of Lagrange's theorem is true for some groups (*e.g.*, all cyclic groups), but it is false in general: given a divisor of the order of the group, there need not be a subgroup with that size.

Example 6.11. The smallest counterexample is A_4 , which has order 12. While A_4 has subgroups of order 1, 2, 3, 4, and 12, it has no subgroup of order 6.

To prove there is no subgroup of order 6, we argue by contradiction. Suppose there is a subgroup of A_4 with order 6, say H . Then $[A_4 : H] = 12/6 = 2$. We will show for each $g \in A_4$ that $g^2 \in H$.

If $g \in H$ then clearly $g^2 \in H$. If $g \notin H$ then gH is a left coset of H different from H (since $g \in gH$ and $g \notin H$), so from $[G : H] = 2$ the only left cosets of H are H and gH . Which one is g^2H ? If $g^2H = gH$ then $g^2 \in gH$, so $g^2 = gh$ for some $h \in H$, and that implies $g = h$, so $g \in H$, but that's a contradiction. Therefore $g^2H = H$, so $g^2 \in H$.

Every 3-cycle (abc) in A_4 is a square: (abc) has order 3, so $(abc) = (abc)^4 = ((abc)^2)^2$. Thus H contains all 3-cycles in A_4 . The 3-cycles are

$$(123), (132), (124), (142), (134), (143), (234), (243)$$

and that is too much since there are 8 of them while $|H| = 6$. Hence H does not exist.

Example 6.12. The next counterexample to the full converse of Lagrange's theorem, in terms of size, is $\mathrm{SL}_2(\mathbf{Z}/(3))$. This group has order 24. It has subgroups of order 1, 2, 3, 4, 6, 8, and 24, but no subgroup of order 12.

To see why there is no subgroup of order 12, assume otherwise. Each subgroup H of order 12 in $\mathrm{SL}_2(\mathbf{Z}/(3))$ has index 2, so just as in the previous example one can show a square in $\mathrm{SL}_2(\mathbf{Z}/(3))$ must lie in H . A tedious count shows there are 10 squares in $\mathrm{SL}_2(\mathbf{Z}/(3))$. Since $|H| = 12$, we don't yet have a contradiction. To reach a contradiction, we use the two squares $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}^2$ and $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}^2$. They must be in H , so their products $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ lie in H . These products turn out not to be squares, so they fill up the available space in H . The inverses of these products also have to lie in H , but an explicit calculation shows the inverses of these two products are matrices not yet taken into account. Thus H is too small, so H does not exist.

Example 6.13. For $n \geq 5$, A_n has no proper subgroup with index less than n ,³ so A_5 has no subgroup with index 2, 3, or 4 even though it has subgroups of order 2, 3 or 4.

Remark 6.14. There is no simple dividing line between those finite groups that satisfy the converse to Lagrange's theorem (for each divisor, there is a subgroup of that order) and those finite groups that do not. Three classes of finite groups that satisfy the converse include abelian groups, dihedral groups, and groups of prime-power order. (These are all special cases of "supersolvable" groups, and all supersolvable groups satisfy the converse of Lagrange's theorem, but some non-supersolvable groups also satisfy the converse of Lagrange's theorem.)

Since we defined the index of a subgroup as the number of its left cosets, presumably we need to introduce a corresponding index for the count of right cosets. However, an extra "right index" concept is not necessary, as we now explain.

Theorem 6.15. *For every subgroup H of G , there are as many left H -cosets as right H -cosets.*

Proof. We will give two proofs, one in the case of finite G and the other in the case of general G .

The "right" analogue of Theorem 5.9 says that each right H -coset is in bijection with H . Therefore, supposing G is finite, running through the proof of Theorem 6.8 with right cosets in place of left cosets shows $|G|/|H|$ is a formula for the number of right H -cosets in G . We already saw this is a formula for the number of left H -cosets, so the number of left and right H -cosets is the same.

Now suppose G is an arbitrary group, possibly infinite. We want to give a bijection between the collections of left and right H -cosets. One's first guess, to send gH to Hg , is *not well-defined*. For instance, taking $G = S_3$ and $H = \{(1), (12)\}$, we have

$$(13)H = (123)H = \{(13), (123)\},$$

but

$$H(13) = \{(13), (132)\}, \quad H(123) = \{(23), (123)\}.$$

So passing from gH to Hg depends on the coset representative g , which means it makes no sense as a function from cosets to cosets. (If you want to remember an example of an attempt to define a function that is in fact *not* well-defined, this is it.)

³See Corollary 8.1 in <https://kconrad.math.uconn.edu/blurbs/grouptheory/Ansimple.pdf>.

The correct way to turn left cosets into right cosets is to use inversion. For a subset $S \subset G$, let $S^{-1} = \{s^{-1} : s \in S\}$. For instance, since subgroups are closed under inversion, check that $H^{-1} = H$ and $(H^{-1})^{-1} = H$. If we invert a left coset gH , we obtain

$$(gH)^{-1} = H^{-1}g^{-1} = Hg^{-1}.$$

Similarly, $(Hg)^{-1} = g^{-1}H$. The function $f(gH) = (gH)^{-1}$ gives a bijection between left and right H -cosets. \square

Theorem 6.15 does not say every right coset is a left coset, but only that the number of each kind of coset is the same. For instance, we saw in Example 5.4 that the different left cosets of $H = \{(1), (12)\}$ in S_3 are

$$\{(1), (12)\}, \{(13), (123)\}, \{(23), (132)\}.$$

The right coset $H(13) = \{(13), (132)\}$ is not the same as a left coset. However, the collection of right H -cosets has 3 members:

$$\{(1), (12)\}, \{(13), (132)\}, \{(23), (123)\}.$$

7. APPLICATIONS OF LAGRANGE'S THEOREM

Lagrange's theorem leads to group-theoretic explanations of some divisibility properties of integers. The idea is this: to prove $a \mid b$, find a group of order b containing a subgroup of order a . After illustrating this in a few cases, we will use Lagrange's theorem to extract information about subgroups of a group.

Theorem 7.1. *Binomial coefficients are integers: for $n \geq 1$ and $0 \leq m \leq n$, the ratio*

$$\binom{n}{m} = \frac{n!}{m!(n-m)!}$$

is an integer.

Proof. The order of S_n is $n!$. We are going to write down a subgroup with order $m!(n-m)!$.

Consider the permutations in S_n that separately permute the subsets $\{1, \dots, m\}$ and $\{m+1, \dots, n\}$:

$$H = \{\sigma \in S_n : \sigma \text{ permutes } \{1, 2, \dots, m\} \text{ and } \{m+1, \dots, n\}\}.$$

The reader can check H is a subgroup, and $|H| = m!(n-m)!$. If the reader is concerned about the degenerate cases $m=0$ and $m=n$, where one of the conditions defining H is an empty condition, note these cases of the theorem are easily checked directly. \square

Theorem 7.2. *For positive integers a and b , the ratios*

$$\frac{(ab)!}{(a!)^b}, \quad \frac{(ab)!}{(a!)^b b!}$$

are integers.

Proof. Write out the integers from 1 to ab in sets of a consecutive integers:

$$1, 2, \dots, a; a+1, \dots, 2a; 2a+1, \dots, 3a; \dots; (b-1)a+1, \dots, ba.$$

There are b of these sets, separated by semi-colons. The permutations of $1, \dots, ba$ that permute each set within itself form a subgroup of S_{ab} and has order $(a!)^b$. Therefore $(a!)^b \mid (ab)!$.

The stronger divisibility relation $(a!)^b b! \mid (ab)!$ can be explained by finding a subgroup of S_{ab} with order $(a!)^b b!$. This is left as an exercise for the reader. Hint: The new subgroup of S_{ab} should contain the one constructed in the previous paragraph. \square

Integrality of the first ratio in the theorem generalizes to $\frac{n!}{k_1!k_2!\cdots k_d!} \in \mathbf{Z}$ when $k_1 + k_2 + \cdots + k_d = n$ with $d \geq 1$ and all k_i in \mathbf{N} .

Theorem 7.3. For $m \geq 1$, let $\varphi(m) = |(\mathbf{Z}/(m))^\times|$ be the number of invertible numbers modulo m . If $m \geq 3$, then $\varphi(m)$ is even.

Here is a table illustrating the evenness of $\varphi(m)$ once $m \geq 3$. Can you find your own explanation?

m	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$\varphi(m)$	1	1	2	2	4	2	6	4	6	4	10	4	12	6	8

Proof. For $m \geq 3$, $\{\pm 1\}$ is a subgroup of $(\mathbf{Z}/(m))^\times$ with order 2, so $2 \mid \varphi(m)$ by Lagrange. \square

Remark 7.4. Curiously, the initial partial sums of the numbers in the above table, added to 1, are often but not always prime. See the table below. Composite values are in red and some primes that are skipped include 17, 31, 37, and 41.

m	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$1 + \sum_{k=1}^m \varphi(k)$	2	3	5	7	11	13	19	23	29	33	43	47	59	65	73

Now we turn to applications of Lagrange's theorem in group theory itself. (The previous application to $\varphi(m)$ is more properly considered an application to number theory than to group theory.) In each application below, pay attention to how Lagrange's theorem is used.

Theorem 7.5. Let G be a finite group and let H and K be subgroups with relatively prime orders. Then $H \cap K = \{e\}$.

Proof. Since $H \cap K$ is a subgroup of both H and K , its order divides both $|H|$ and $|K|$ by Lagrange. Therefore, by relative primality of these orders, $|H \cap K| = 1$, so $H \cap K = \{e\}$. \square

Theorem 7.6. Let G be a finite group. If g has order n , then $n \mid |G|$. In particular, $g^{|G|} = e$ for every $g \in G$.

Proof. Let $H = \langle g \rangle$ be the subgroup generated by g . The order of H is exactly the order of g , so $|H| = n$. By Lagrange, $n \mid |G|$. Since $g^n = e$ and $n \mid |G|$, we get $g^{|G|} = e$. \square

This theorem can be proved for abelian groups by a simpler method: see Corollary 4.3 in <https://kconrad.math.uconn.edu/blurbs/grouptheory/order.pdf>. In the abelian case, we can first prove $g^{|G|} = e$ and then deduce that the order of each element is a factor of $|G|$. In Theorem 7.6, which allows groups to be nonabelian, the proof first shows each element has order dividing $|G|$ and then deduces that $g^{|G|} = e$ for each g in G .

Corollary 7.7. If G is a finite group and k is an integer relatively prime to $|G|$, then the k -th power function on G is invertible.

This is not saying the k -th power function is multiplicative on G , but only that it is a bijection.

Proof. By Bezout, when $(k, |G|) = 1$ we can write $k\ell + |G|m = 1$ for some integers ℓ and m . Therefore

$$g = g^1 = g^{k\ell} (g^{|G|})^m = g^{k\ell}$$

since $g^{|G|} = 1$. So we see how to invert the k -th power function: raise to the power ℓ , where $k\ell \equiv 1 \pmod{|G|}$. \square

Example 7.8. When $G = D_5$, which has order 10, let's raise all the elements to the third power. The result is a rearrangement of the elements of the group. See the table below. (Reflections don't change since an odd power of a reflection is itself.)

g	1	r	r^2	r^3	r^4	s	rs	r^2s	r^3s	r^4s
g^3	1	r^3	r	r^4	r^2	s	rs	r^2s	r^3s	r^4s

Since $3 \cdot 7 \equiv 1 \pmod{10}$, the seventh power is the inverse of the third power: $(g^3)^7 = g^{21} = g \cdot (g^{10})^2 = g$. For example, $(r^4)^3 = r^{12} = r^2$ and $(r^2)^7 = r^{14} = r^4$.

What about the converse of Corollary 7.7? That is, is the k -th power function on G invertible *only* when k is relatively prime to $|G|$? This can be answered using Cauchy's theorem, but we don't do that here.

Corollary 7.9. *Every group of prime order is cyclic, and in fact each non-identity element is a generator.*

Proof. Let G be the group, with $p = |G|$. Pick a non-identity element g from G . By Theorem 7.6, the order of g divides p and is greater than 1, so g has order p . Therefore $|\langle g \rangle| = p$, so $\langle g \rangle = G$. \square

Remark 7.10. Be careful not to mis-apply Corollary 7.9. While it tells us that the additive group $\mathbf{Z}/(p)$ is cyclic (each non-zero number modulo p is an additive generator), it does not tell us anything about the nature of the multiplicative group $(\mathbf{Z}/(p))^\times$, which has order $p - 1$. The group $(\mathbf{Z}/(p))^\times$ is cyclic for prime p , but the proof of that is not as simple as for $\mathbf{Z}/(p)$. It certainly does not come from Corollary 7.9, since $(\mathbf{Z}/(p))^\times$ does not have prime order (when $p > 3$).

Corollary 7.11. *Let p and q be primes. Each non-trivial proper subgroup of a group of order pq is cyclic.*

Proof. A nontrivial proper subgroup will have order equal to a non-trivial factor of pq , which is either p or q . In either case, the subgroup has prime order and therefore is cyclic by Corollary 7.9. \square

For instance, if we want to find all the subgroups of a group of order 6 (or 15 or 21...), we can compute the cyclic group generated by each element. Every non-trivial proper subgroup will arise in this way, and we then throw in the trivial group and the whole group to complete the list.

Here is an application of Lagrange's theorem to a characterization of cyclic groups from combinatorial information about their subgroups.

Theorem 7.12. *A finite group that has at most one subgroup of each size is cyclic.*

Proof. I learned this proof from F. Ladisch.

Let G be a group with at most one subgroup of each size, and let $n = |G|$. We will count the elements of G by *collecting together all elements that generate the same subgroup*. That is, for each cyclic subgroup H we will collect together all of its generators and then add up

the number of these generators as H varies. This accounts for each element of G exactly once (each element generates some cyclic subgroup), so

$$n = \sum_{\text{cyc. subgp. } H} |\{\text{generators of } H\}|.$$

Every subgroup of G has order dividing n , by Lagrange's theorem, so we can sum over cyclic subgroups of G by summing over cyclic subgroups of order d for each d dividing n :

$$n = \sum_{d|n} \sum_{\text{cyc. } H, |H|=d} |\{\text{generators of } H\}|.$$

All cyclic groups of order d have the same number of generators: if $\langle h \rangle$ has order d then each element of $\langle h \rangle$ is h^a for a unique a between 1 and d , and h^a generates $\langle h \rangle$ if and only if $(a, d) = 1$. Thus the number of generators in a cyclic group of order d is $|\{1 \leq a \leq d : (a, d) = 1\}| = \varphi(d)$, so

$$(7.1) \quad n = \sum_{d|n} \sum_{\text{cyc. } H, |H|=d} \varphi(d) = \sum_{d|n} \varphi(d) |\{H \subset G : |H| = d, H \text{ cyclic}\}|.$$

This equation is valid for *every* finite group G of order n .

In the particular group $\mathbf{Z}/(n)$ of order n , which is cyclic, there is a unique cyclic subgroup of every order dividing n , so (7.1) for $G = \mathbf{Z}/(n)$ becomes

$$(7.2) \quad n = \sum_{d|n} \varphi(d).$$

If G is a group of order n and it has at most one subgroup of order d for each d dividing n , then $|\{H \subset G : |H| = d, H \text{ cyclic}\}| = 0$ or 1 for all d dividing n , so

$$(7.3) \quad n = \sum_{d|n} \varphi(d) |\{H \subset G : |H| = d, H \text{ cyclic}\}| \leq \sum_{d|n} \varphi(d) \stackrel{(7.2)}{=} n.$$

Therefore $|\{H \subset G : |H| = d, H \text{ cyclic}\}| = 1$ for all d dividing n : if the count were ever 0 then the inequality in (7.3) would be strict and hence $n < n$, a contradiction. Since G has a cyclic subgroup of order d for each d dividing n , taking $d = n$ shows G is cyclic. \square

The only subgroups used in this proof are cyclic, so we proved something slightly stronger than what was stated in the theorem: a finite group that has at most one *cyclic* subgroup of each size is cyclic.

Corollary 7.13. *Let G be a finite group such that, for each d dividing $|G|$, the equation $x^d = 1$ in G has at most d solutions. Then G is cyclic.*

Proof. We will show G has at most one subgroup of each size. If H is a subgroup of G with order d , then every element of H satisfies the equation $x^d = 1$ and thus, by hypothesis, H is the complete set of solutions in G to the equation $x^d = 1$. This shows there is at most one subgroup of order d : if H and H' are both subgroups of order d then H and H' both equal $\{g \in G : g^d = 1\}$, so $H = H'$. \square

We conclude our list of applications of Lagrange's theorem with a result that stands in some sense "dual" to Theorem 7.5: instead of the subgroups having relatively prime orders, and getting information about the size of the intersection, we look at what happens when the subgroups have relatively prime index. What can be said about the index of their intersection?

Theorem 7.14. *Let G be a finite group, with subgroups H and K . Set $m = [G : H]$ and $n = [G : K]$. Then*

$$[m, n] \leq [G : H \cap K] \leq mn.$$

In particular, if m and n are relatively prime, then $[G : H \cap K] = mn = [G : H][G : K]$.

Proof. Since $H \cap K \subset H \subset G$ and $H \cap K \subset K \subset G$, Theorem 6.9 tells us

$$[G : H \cap K] = [G : H][H : H \cap K] = [G : K][K : H \cap K].$$

Thus m and n each divide $[G : H \cap K]$, so their least common multiple divides $[G : H \cap K]$ as well.

Now we want to show $[G : H \cap K] \leq mn$. Writing this as

$$[G : H][H : H \cap K] \leq [G : H][G : K],$$

our desired inequality is the same as

$$(7.4) \quad [H : H \cap K] \leq [G : K].$$

The number $[G : K]$ counts how many left K -cosets are in G . Every left K -coset has the form gK for some $g \in G$. Among these, some cosets contain elements of H . How many? We will show there are $[H : H \cap K]$ such cosets, and thus we obtain the inequality (7.4).

The point is, for h_1 and h_2 in H , that

$$(7.5) \quad h_1K = h_2K \iff h_1(H \cap K) = h_2(H \cap K).$$

(If this is true, then it tells us the number of left K -cosets of G represented by an element of H is the same as the number of left $(H \cap K)$ -cosets in H , since $H \cap K$ is a subgroup of H .) Well, on the left side of (7.5), there is such equality exactly when $h_1 = h_2k$ for some $k \in K$. Then $k = h_2^{-1}h_1$ lies in H as well, so $k \in H \cap K$. But then the equation $h_1 = h_2k$ tells us $h_1(H \cap K) = h_2(H \cap K)$. The reverse implication in (7.5) is left as an exercise for the interested reader. \square

8. LEFT AND RIGHT COSET SPACES

The usefulness of modular arithmetic (in number theory, cryptography, *etc.*) suggests the possibility of carrying it over from \mathbf{Z} to other groups. After all, since congruence classes modulo $m\mathbf{Z}$ are just a special instance of a coset decomposition, why not think about (left) cosets for subgroups of groups other than \mathbf{Z} as a generalization of $\mathbf{Z}/(m)$?

Definition 8.1. Let G be a group and H be a subgroup. The *left coset space* $G/H := \{gH : g \in G\}$ is the set consisting of the left H -cosets in G .

A left coset space is a set whose elements are the left cosets of a subgroup. It is a “set of sets,” as the following examples illustrate.

Example 8.2. In the group S_3 , let $H = \{(1), (12)\}$. By (5.2), there are 3 left H -cosets:

$$\begin{aligned} S_3/H &= \{H, (13)H, (23)H\} \\ &= \{\{(1), (12)\}, \{(13), (123)\}, \{(23), (132)\}\}. \end{aligned}$$

Example 8.3. In $\text{Aff}^+(\mathbf{R})$, with $K = \left\{ \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix} : x > 0 \right\}$, each left K -coset contains one matrix of the form $\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$, so

$$\text{Aff}^+(\mathbf{R})/K = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} K : b \in \mathbf{R} \right\}.$$

Example 8.4. In \mathbf{R} , every \mathbf{Z} -coset is represented by a real number in $[0, 1)$, so (5.5) tells us

$$\mathbf{R}/\mathbf{Z} = \{x + \mathbf{Z} : 0 \leq x < 1\}.$$

Can we make G/H into a group using the group operation from G , in the same way that $\mathbf{Z}/(m)$ inherits an additive group structure from that of \mathbf{Z} ? The natural idea would be to multiply two left cosets by multiplying two representatives:

$$g_1H \cdot g_2H := g_1g_2H.$$

Unfortunately, such an operation does not always make sense. That is, the coset on the right can change if we change the coset representatives on the left without changing the cosets themselves.

Example 8.5. Take $G = S_3$ and $H = \{(1), (12)\}$. We have

$$(13)H = (123)H = \{(13), (123)\}, \quad (23)H = (132)H = \{(23), (132)\},$$

but

$$(13)(23)H = (132)H, \quad (123)(132)H = H,$$

and $(132)H \neq H$, so changing the coset representatives changed the ‘product’ coset.

In Example 8.5, multiplication of left cosets is *not well-defined*: it depends on the choice of coset representative, and thus is not an honest operation at the level of cosets.

When G is abelian, this kind of difficulty does not arise: if $g_1H = g'_1H$ and $g_2H = g'_2H$, then $g_1g_2H = g'_1g'_2H$. One can proceed and turn G/H into a group, just as $\mathbf{Z}/(m)$ is a group for addition. When G is non-abelian (the far more typical case), there are special kinds of subgroups, called *normal* subgroups, for which multiplication of left cosets is a well-defined operation in terms of representatives, and one can make G/H into a group by multiplying coset representatives.

Why should we care about trying to make G/H into a group? Here are three reasons:

- The usefulness of modular arithmetic in $\mathbf{Z}/(m)$ suggests having an analogous construction for other groups has got to be worthwhile, even if it is not yet clear what we might be able to do with the construction. (Note: While we can both add and multiply in $\mathbf{Z}/(m)$, we only have in mind that G/H could be a group, inheriting the one operation on G , analogous just to addition in $\mathbf{Z}/(m)$.)
- Having groups of the form G/H is a method of constructing new groups out of old ones (by replacing G with G/H). This could allow us to find alternate models for certain kinds of groups, which might be more convenient to use than other models.
- When G is finite and H is a non-trivial proper subgroup of G , both H and G/H have order less than the order of G . If G/H is a group, then we have two groups related to G that have order less than $|G|$. This turns out to be very useful in proving theorems about finite groups by induction on the order of the group.

Remark 8.6. In addition to the left coset space G/H , one can also consider the right H -coset space $H \backslash G := \{Hg : g \in G\}$.

9. DOUBLE COSETS

A coset can be viewed as the result of multiplying an element on one side by a subgroup. Allowing multiplying on both sides (by possibly different subgroups) leads to the idea of a double coset.

Definition 9.1. Let G be a group and H and K be subgroups. A set of the form

$$HgK = \{h g k : h \in H, k \in K\}$$

is called an (H, K) *double coset*, or simply a double coset if H and K are understood.

While special instances of double cosets appeared in the work of Cauchy, their systematic consideration in group theory is due to Frobenius.

In this section we will look at some examples of double cosets and see how they are similar to and different from left and right cosets.

Example 9.2. Let's look at double cosets in S_3 , with $H = \{(1), (12)\}$ and $K = \{(1), (13)\}$. The double coset of (1) is

$$H(1)K = HK = \{hk : h \in H, k \in K\} = \{(1), (12), (13), (132)\}$$

and the double coset of (23) is

$$\begin{aligned} H(23)K &= \{(1)(23)(1), (1)(23)(13), (12)(23)(1), (12)(23)(13)\} \\ &= \{(23), (123)\}. \end{aligned}$$

Unlike a coset, we now see that a double coset does not have to have order dividing the order of the group (4 does not divide 6). Moreover, different double cosets for the same pair of subgroups can have different orders (such as 4 and 2).

The reader can compute the (H, K) -double cosets of the other four elements of S_3 and find only the two examples above repeating. (Or appeal to Theorem 9.7 below.)

Example 9.3. Consider the group S_3 again, but now use $H = K = \{(1), (12)\}$. There are only two different double cosets:

$$H(1)H = HH = \{(1), (12)\}, \quad H(13)H = \{(13), (23), (123), (132)\}.$$

Example 9.4. Take $G = D_4 = \langle r, s \rangle$ and $H = K = \{1, s\}$. The different (H, H) double cosets are

$$HH = H = \{1, s\}, \quad HrH = \{r, rs, r^3s, r^3\}, \quad Hr^2H = \{r^2, r^2s\}.$$

This example shows the number of double cosets for a pair of subgroups need not divide the order of the group (3 does not divide 8). This is a contrast to what happens with left and right cosets of a subgroup.

Example 9.5. An ordinary coset is a special kind of double coset, where one of the subgroups is trivial: $Hg\{e\} = Hg$ and $\{e\}gK = gK$.

Example 9.6. If G is abelian, then the product set HK is a subgroup of G and an (H, K) double coset is just an ordinary coset of the subgroup HK .

Theorem 9.7. *Fix two subgroups H and K of the group G . Every element of G lies in some (H, K) double coset, and two (H, K) double cosets that overlap are equal. Equivalently, different (H, K) double cosets are disjoint and they collectively cover the whole group.*

Proof. Clearly $g \in HgK$: use $h = e$ and $k = e$. Thus, every element of G is in some (H, K) double coset.

Now assume HgK and $Hg'K$ overlap:

$$(9.1) \quad h g k = h' g' k',$$

where h and h' belong to H and k and k' belong to K . We want to prove $HgK = Hg'K$. By (9.1) $g = h^{-1}h'g'k'k^{-1}$, so g lies in $Hg'K$. For h'' in H and k'' in K ,

$$h''gk'' = h''h^{-1}h'g'k'k^{-1}k'' \in Hg'K.$$

Letting h'' and k'' vary, we obtain $HgK \subset Hg'K$. The reverse inclusion is proved in the same way, so $HgK = Hg'K$. \square

A generalization of left and right coset spaces in Section 8 is a double coset space

$$H \backslash G / K = \{HgK : g \in G\}.$$

This is a left or right coset space when H or K is trivial. Inverting elements turns G/H into $H \backslash G$. Similarly, inverting turns $H \backslash G / K$ into $K \backslash G / H$.

We noted in Example 9.4 that the number of (H, K) double cosets does not have to divide the order of the group. However, Frobenius found a formula involving double coset spaces that does generalize the “index formulas” $|G| = |H||G/H| = |H||H \backslash G|$, as follows.

Theorem 9.8. *Let G be a finite group and H and K be subgroups. Then*

$$|\{(h, g, k) \in H \times G \times K : h g k = g\}| = |H||K||H \backslash G / K|.$$

When H or K is trivial, this reduces to the index formula for the other subgroup.

10. HISTORY

The first use of cosets in a group-theoretic setting was by Euler in the mid-1700s, when he proved [2, Theorems 10–14] for prime p and $a \not\equiv 0 \pmod{p}$ that $a^{p-1} \equiv 1 \pmod{p}$ by reasoning as in the proof of Theorem 7.6. More specifically, he showed the subgroup $H = \langle a \pmod{p} \rangle$ of the group $G = (\mathbf{Z}/(p))^\times$ has order equal to the order of $a \pmod{p}$ and then proved (6.2) in this case by using the cosets of H in G . Later [3, Theorems 8–11], [4, Sect. 195–213] he used cosets to show that if $(a, m) = 1$ then $a^{\varphi(m)} \equiv 1 \pmod{m}$.

While group theory as its own branch of math took form in the 19th century, the term “coset” was introduced in a paper by G. A. Miller [6] in 1910, written as co-set. The prefix “co-” has its standard meaning of “together with”, as in the word coauthor. Before this paper, Burnside’s book *Theory of Groups of Finite Order* from 1897 used right cosets to prove Lagrange’s theorem [1, pp. 21–22], but it had no special terminology for cosets. Weber’s *Lehrbuch der Algebra* from 1896 did have a name (in German) for cosets: Nebengruppen (“nearby groups”) [7, p. 8]. Strictly speaking, Weber used this label for cosets of a subgroup other than the subgroup itself. In German today, the term for coset is Nebenklasse.

The label “Lagrange’s theorem” also became standard in the 20th century. It appears in the 19th century literature [5, p. 273], but not in Burnside [1, p. 21] or Weber [7, p. 8].

REFERENCES

- [1] W. Burnside, “Theory of Groups of Finite Order,” Cambridge Univ. Press, Cambridge, 1897. URL <https://archive.org/details/in.ernet.dli.2015.168629/mode/2up>.
- [2] L. Euler, Theoremata circa residua ex divisione potestatum relictia, *Novi Commentarii Academiae Scientiarum Imperialis Petropolitanae* **7** (1761), 49–82. URL <https://scholarlycommons.pacific.edu/euler-works/262/>.
- [3] L. Euler, Theoremata arithmetica nova methodo demonstrata, *Novi Commentarii Academiae Scientiarum Imperialis Petropolitanae* **8** (1763), 74–104. URL <https://scholarlycommons.pacific.edu/euler-works/271/>. German translation at <https://arxiv.org/abs/1203.1993>.
- [4] L. Euler, Tractatus de numerorum doctrina capita sedecim, quae supersunt, *Commentationes Arithmeticae* **2** (1849), 503–575. URL <https://scholarlycommons.pacific.edu/euler-works/792/>.

- [5] G. Frobenius, Ueber die Congruenz nach einem aus zwei endlichen Gruppen gebildeten Doppelmodul, *Journal für die reine und angewandte Mathematik* **101** (1887), 273–299. URL <https://eudml.org/doc/148695>.
- [6] G. A. Miller, On a method due to Galois, *Quarterly Journal of Mathematics* **41** (1910), 382–384. URL <https://babel.hathitrust.org/cgi/pt?id=inu.30000050138258&seq=390>.
- [7] H. Weber, “Lehrbuch der Algebra,” Band II, F. Vieweg und Sohn, Braunschweig, 1896. URL <https://archive.org/details/lehrbuchderalgeb02weberich/page/n7/mode/2up>.