

## A “UNIVERSAL” DIVISIBILITY TEST

KEITH CONRAD

The tests for divisibility by 3, 9, and 11 all have a similar flavor: for a positive integer

$$(1) \quad n = a_d 10^d + a_{d-1} 10^{d-1} + \cdots + a_1 10 + a_0,$$

where  $0 \leq a_i \leq 9$ , we have

$$\begin{aligned} n \equiv 0 \pmod{3} &\iff a_d + a_{d-1} + a_{d-2} + \cdots + a_0 \equiv 0 \pmod{3}, \\ n \equiv 0 \pmod{9} &\iff a_d + a_{d-1} + a_{d-2} + \cdots + a_0 \equiv 0 \pmod{9}, \\ n \equiv 0 \pmod{11} &\iff a_d - a_{d-1} + a_{d-2} - \cdots + (-1)^d a_0 \equiv 0 \pmod{11}. \end{aligned}$$

These tests are convenient to use because the sum of the digits of  $n$  and the alternating sum of the digits of  $n$  are much smaller than  $n$ , so we can turn the divisibility problem for  $n$  into a divisibility problem for a smaller number. Moreover, we can iterate the test again and again until we are left with a very small number to test.

These three tests generalize to a test for divisibility by *any* integer  $m$  relatively prime to 10 (that is,  $m$  is not a multiple of 2 or 5). So we will get, for instance, divisibility tests by 7, 13, and 29. The general test will involve the operation of taking off the units' digit of a positive integer, *e.g.*, turning 1634 into 163 or 78325 into 7832. For  $n \geq 1$ , let  $n'$  be the number that we get after taking off the units' digit of  $n$ . So if  $n$  is written as in (1),

$$(2) \quad n' = a_d 10^{d-1} + a_{d-1} 10^{d-2} + \cdots + a_1 = \frac{n - a_0}{10}.$$

In (2) we removed the digit  $a_0$  and shifted all the other digits into the next lower position ( $a_1$  fills the position previously taken by  $a_0$ , and so on).

Here is the universal divisibility test.

**Theorem 1.** *When  $(m, 10) = 1$ , choose  $b$  so that  $10b \equiv 1 \pmod{m}$ . Then*

$$n \equiv 0 \pmod{m} \iff n' + ba_0 \equiv 0 \pmod{m}.$$

We will look at a number of examples of this before we discuss the proof.

**Example 2.** Take  $m = 7$ . Then  $10 \cdot 5 \equiv 1 \pmod{7}$ , so

$$(3) \quad n \equiv 0 \pmod{7} \iff n' + 5a_0 \equiv 0 \pmod{7}.$$

Let's try  $n = 11382$ . We have  $n' = 1138$  and  $n' + 5a_0 = 1138 + 5 \cdot 2 = 1148$ , so  $7|n$  if and only if  $7|1148$ . Since 1148 is still big, we apply the test again to 1148:  $114 + 5 \cdot 8 = 114 + 40 = 154$ , so  $7|1148$  if and only if  $7|154$ . Then we replace 154 with  $15 + 5 \cdot 4 = 15 + 20 = 35$ , which *is* divisible by 7. Thus the original number 11382 is divisible by 7 (because the test is an “if and only if” criterion, so it works in both directions). Explicitly,

$$n = 11382 = 7 \cdot 1626.$$

Let's summarize our successive computations in the following way:

$$11382 \rightsquigarrow 1138 + 5 \cdot 2 = 1148 \rightsquigarrow 114 + 5 \cdot 8 = 154 \rightsquigarrow 15 + 5 \cdot 4 = 35.$$

Any  $b$  fitting  $10b \equiv 1 \pmod{7}$  can be used in place of 5 in this test. Since  $10(-2) \equiv 1 \pmod{7}$ , for instance, we also get a test for divisibility by 7 as

$$(4) \quad n \equiv 0 \pmod{7} \iff n' - 2a_0 \equiv 0 \pmod{7}.$$

This is more convenient to use than (3) since  $-2$  is smaller in magnitude than 5. Of course (3) and (4) are the same test, since  $5 \equiv -2 \pmod{7}$ , but the integers they lead to are different. Let's apply (4) to 11382. The successive numbers we get now are

$$11382 \rightsquigarrow 1138 - 2 \cdot 2 = 1134 \rightsquigarrow 113 - 2 \cdot 4 = 105 \rightsquigarrow 10 - 2 \cdot 5 = 0,$$

which is divisible by 7, so the original number 11382 is divisible by 7.

**Example 3.** Take  $m = 13$ . Then  $10 \cdot 4 \equiv 1 \pmod{13}$ , so

$$n \equiv 0 \pmod{13} \iff n' + 4a_0 \equiv 0 \pmod{13}.$$

Taking  $n = 11382$  again, the successive numbers under the operation  $n \rightsquigarrow n' + 4a_0$  are

$$11382 \rightsquigarrow 1138 + 4 \cdot 2 = 1146 \rightsquigarrow 114 + 4 \cdot 6 = 138 \rightsquigarrow 13 + 4 \cdot 8 = 45 \rightsquigarrow 4 + 4 \cdot 5 = 24,$$

which is *not* divisible by 13, so 11382 is not divisible by 13. (We didn't have to stop there:  $24 \rightsquigarrow 2 + 4 \cdot 4 = 18$ , which is not divisible by 13.) Trying now  $n = 78325$ , we compute

$$78325 \rightsquigarrow 7832 + 4 \cdot 5 = 7852 \rightsquigarrow 785 + 4 \cdot 2 = 793 \rightsquigarrow 79 + 4 \cdot 3 = 91 \rightsquigarrow 9 + 4 = 13,$$

so 78325 is divisible by 13. Explicitly,

$$78325 = 13 \cdot 6025.$$

For  $m < 50$  with  $(10, m) = 1$ , Table 1 below lists the inverse of  $10 \pmod{m}$  in the second column, using the representative that is smallest in absolute value (so for  $m = 7$  we choose  $-2$  rather than 5).

**Example 4.** From Table 1,

$$n \equiv 0 \pmod{17} \iff n' - 5a_0 \equiv 0 \pmod{17},$$

$$n \equiv 0 \pmod{19} \iff n' + 2a_0 \equiv 0 \pmod{19},$$

$$n \equiv 0 \pmod{21} \iff n' - 2a_0 \equiv 0 \pmod{21},$$

$$n \equiv 0 \pmod{23} \iff n' + 7a_0 \equiv 0 \pmod{23},$$

$$n \equiv 0 \pmod{27} \iff n' - 8a_0 \equiv 0 \pmod{27},$$

and

$$n \equiv 0 \pmod{29} \iff n' + 3a_0 \equiv 0 \pmod{29}.$$

Let's see if 1634 is divisible by 29. The operation is  $n \rightsquigarrow n' + 3a_0$  in this case, and

$$1634 \rightsquigarrow 163 + 3 \cdot 4 = 175 \rightsquigarrow 17 + 3 \cdot 5 = 32,$$

which is not a multiple of 29, so 1634 is not divisible by 29. Now trying 13108, we get

$$13108 \rightsquigarrow 1310 + 3 \cdot 8 = 1334 \rightsquigarrow 133 + 3 \cdot 4 = 145 \rightsquigarrow 14 + 3 \cdot 5 = 29,$$

which is divisible by 29, so 13108 is divisible by 29. Explicitly,

$$13108 = 29 \cdot 452.$$

Now that we see how Theorem 1 works in practice, let's prove it. The proof will be very short! It depends on writing  $n$  as  $10n' + a_0$  and doing one multiplication mod  $m$ .

$m$	$b$
3	1
7	-2
9	1
11	-1
13	4
17	-5
19	2
21	-2
23	7
27	-8
29	3
31	-3
33	10
37	-11
39	4
41	-4
43	13
47	-14
49	5

TABLE 1. A solution to  $10b \equiv 1 \pmod{m}$ 

*Proof.* Since  $n = 10n' + a_0$ ,

$$n \equiv 0 \pmod{m} \iff 10n' + a_0 \equiv 0 \pmod{m}.$$

Since  $10 \pmod{m}$  is invertible, with inverse  $b$ ,

$$\begin{aligned} 10n' + a_0 \equiv 0 \pmod{m} &\iff b(10n' + a_0) \equiv 0 \pmod{m} \\ &\iff n' + ba_0 \equiv 0 \pmod{m}. \end{aligned}$$

□

All that really happened in the proof is that we divided by 10 working modulo  $m$ . If we allow ourselves to use ordinary fractional notation,  $10n' + a_0 \equiv 0 \pmod{m}$  if and only if  $n' + a_0/10 \equiv 0 \pmod{m}$  and the legal form of  $1/10 \pmod{m}$  is  $b \pmod{m}$  since  $10b \equiv 1 \pmod{m}$ .

Although we said at the start that the divisibility test in Theorem 1 generalizes the divisibility tests for 3, 9, and 11, which involve adding (or alternately adding and subtracting) all the digits of a number, the usual tests for 3, 9, and 11 don't actually look like the test in Theorem 1. So let's see how Theorem 1 implies the usual tests for 3, 9, and 11. Looking at Table 1, where  $b = 1$  for  $m = 3$  and 9, and  $b = -1$  for  $m = 11$ , Theorem 1 says

$$\begin{aligned} n \equiv 0 \pmod{3} &\iff n' + a_0 \equiv 0 \pmod{3}, \\ n \equiv 0 \pmod{9} &\iff n' + a_0 \equiv 0 \pmod{9}, \\ n \equiv 0 \pmod{11} &\iff n' - a_0 \equiv 0 \pmod{11}. \end{aligned}$$

Since  $10 \equiv 1 \pmod{3}$ , by (2)

$$n' \equiv a_d + a_{d-1} + \cdots + a_1 \pmod{3},$$

so

$$n' + a_0 \equiv a_d + a_{d-1} + \cdots + a_1 + a_0 \pmod{3}.$$

Therefore the test for divisibility by 3 in Theorem 1 is the same as

$$n \equiv 0 \pmod{3} \iff a_d + a_{d-1} + \cdots + a_1 + a_0 \equiv 0 \pmod{3},$$

which is the usual test for divisibility by 3. Since  $10 \equiv 1 \pmod{9}$ , Theorem 1 implies the usual test for divisibility by 9 in the same way. As for 11, since  $10 \equiv -1 \pmod{11}$  we have

$$n' \equiv a_d(-1)^{d-1} + a_{d-1}(-1)^{d-2} + \cdots + a_1 \pmod{11},$$

so

$$n' - a_0 \equiv a_d(-1)^{d-1} + a_{d-1}(-1)^{d-2} + \cdots + a_1 - a_0 \pmod{11}.$$

Therefore Theorem 1 says

$$\begin{aligned} n \equiv 0 \pmod{11} &\iff n' - a_0 \equiv 0 \pmod{11} \\ &\iff a_d(-1)^{d-1} + a_{d-1}(-1)^{d-2} + \cdots + a_1 - a_0 \pmod{11} \\ &\iff (-1)^{d-1}(a_d - a_{d-1} + \cdots + (-1)^{d-1}a_1 + (-1)^d a_0) \equiv 0 \pmod{11} \\ &\iff a_d - a_{d-1} + \cdots + (-1)^{d-1}a_1 + (-1)^d a_0 \equiv 0 \pmod{11}, \end{aligned}$$

which is the usual for divisibility by 11.

**Remark 5.** If we try out the universal divisibility test for  $m$  on a number that is too small (relative to  $m$ ), we may produce larger numbers in the recursion. For example, take  $m = 13$  (and  $b = 4$ ). Testing for divisibility of 28 by 13, we get

$$28 \rightsquigarrow 2 + 4 \cdot 8 = 34 \rightsquigarrow 3 + 4 \cdot 4 = 19 \rightsquigarrow 1 + 4 \cdot 9 = 37 \rightsquigarrow 3 + 4 \cdot 7 = 31 \rightsquigarrow 3 + 4 \cdot 1 = 7,$$

which is not divisible by 13 so 28 isn't divisible by 13 either. Notice the sequence went up and down a couple of times before getting very small.

It can also happen that the recursion enters a loop. For example, if we want to test 351 for divisibility by 13 then we get

$$351 \rightsquigarrow 35 + 4 \cdot 1 = 39 \rightsquigarrow 3 + 4 \cdot 9 = 39 \rightsquigarrow 39 \rightsquigarrow 39 \rightsquigarrow \dots$$

It can be shown that the “universal” test for divisibility by  $m$  will lead to rising numbers or a loop only at a stage where the numbers are small relative to  $m$  (of size less than  $10m$ , in fact), at which point you could just stop and do a direct divisibility check.